







# Quanti-CultiControl

The Quanti-CultiControl are lyophilized, quantitative microorganism preparations to be used in industrial laboratories for Quality Control purposes. Processed as directed, these preparations provide a challenge of <100 CFU per 0.1 mL, which is a concentration usually suitable for several pharmaceutical applications, including growth promotion testing of culture media to be employed in sterility testing. These microorganism preparations are derived from the American Type Culture Collection (ATCC®) or other authentic reference culture collection.

The Quanti-CultiControl are suitable for the Growth Promotion Test recommended by the international Pharmacopoeiae, without need of intermediate dilution steps, since each rehydrated suspension generates less than 100 UFC for 0.1 mL inoculum. Each suspension (1 mL) allows 10 inocula.

Liofilchem® and the Liofilchem company logo are registered trademarks of LIOFILCHEM s.r.l.

 <p><b>1.</b></p>	 <p><b>2.</b></p>
<p>Remove the vial of lyophilized pellet from the refrigerator (2-8 °C). Allow the materials to equilibrate to room temperature (about 30 minutes) before opening the vial.</p>	<p>While the pellets are equilibrating, prewarm the hydrating fluid to 34-38°C (at least 30 minutes).</p>
 <p><b>3.</b></p>	 <p><b>4.</b></p>
<p>With a sterile forceps, transfer the pellet into the 1 mL vial of hydrating fluid. Do not remove the desiccant from vial. One pellet must be used to obtain the challenge concentration of 10–100 CFU per 0.1 mL on non-selective media.</p>	<p>Immediately recap the vial with the hydrated material and place into a 34-38 °C incubator for 30 minutes to ensure complete hydration.</p>
 <p><b>5.</b></p>	 <p><b>6.</b></p>
<p>Immediately following incubation, vortex the hydrated material until pellets have completely dissolved and suspension is homogeneous.</p>	<p>With a sterile pipette, transfer 0.1 mL of the hydrated suspension to the material being challenged (0.1 mL contains 10–100 CFU). Remaining suspension can be refrigerated and used for up to 8 hours (no warming time required), with the exception of catalog number 89505 which must be used within 30 minutes.</p>
<p><b>7.</b> Proceed with the challenge procedure according to laboratory protocol. Discard any remaining hydrated material in accordance with the laboratory protocol for disposal of biohazard materials.</p>	



**Liofilchem s.r.l.**  
Roseto degli Abruzzi, Italy  
[www.liofilchem.com](http://www.liofilchem.com)



Microbiology Products since 1983

ref. 6550217  
Rev. 5 - 1.3.2021  
© Liofilchem 2021  
printed in Italy

# Quanti-CultiControl™ freeze-dried microorganisms

**Packaging:** 1 vial containing 1 pellet + 1 rehydration fluid vial

**Quantitative:** <100 CFU / 0.1 mL inoculum

**Applications:** Growth Promotion Testing, Microbial Limits Testing, Microbial Enumeration Testing

Description	Ref.	notes
<i>Aspergillus brasiliensis</i> derived from ATCC® 16404™*	89501	
<i>Bacillus cereus</i> derived from ATCC® 10876™*	89502	
<i>Bacillus subtilis</i> subsp. spizizenii derived from ATCC® 6633™*	89503	
<i>Bacteroides fragilis</i> derived from ATCC® 25285™*	89505	
<i>Bifidobacterium animalis</i> subsp. animalis derived from ATCC® 25527™*	89539	
<i>Brevundimonas diminuta</i> derived from ATCC® 19146™*	89506	
<i>Burkholderia cepacia</i> derived from ATCC® 25416™*	89507	
<i>Candida albicans</i> derived from ATCC® 10231™*	89508	
<i>Candida albicans</i> derived from ATCC® 2091™*	89510	
<i>Clostridium sporogenes</i> derived from ATCC® 11437™*	89513	
<i>Clostridium sporogenes</i> derived from ATCC® 19404™*	89514	
<i>Enterobacter aerogenes</i> derived from ATCC® 13048™*	89516	
<i>Enterococcus faecalis</i> derived from ATCC® 29212™*	89517	
<i>Enterococcus faecalis</i> derived from ATCC® 7080™*	89518	
<i>Escherichia coli</i> derived from ATCC® 8739™*	89519	
<i>Geobacillus stearothermophilus</i> derived from ATCC® 12980™*	89521	
<i>Geobacillus stearothermophilus</i> derived from ATCC® 7953™*	89522	
<i>Kocuria rhizophila</i> derived from ATCC® 9341™*	89523	
<i>Lactobacillus fermentum</i> derived from ATCC® 9338™*	89524	
<i>Listeria monocytogenes</i> derived from ATCC® 19115™*	89525	serotype 4b
<i>Micrococcus luteus</i> derived from ATCC® 4698™*	89526	
<i>Pseudomonas aeruginosa</i> derived from ATCC® 27853™*	89527	
<i>Pseudomonas aeruginosa</i> derived from ATCC® 9027™*	89528	
<i>Salmonella enterica</i> subsp. enterica serovar Typhimurium derived from ATCC® 14028™*	89531	
<i>Salmonella enterica</i> subsp. enterica serovar Typhimurium derived from ATCC® 13311™*	89530	
<i>Staphylococcus aureus</i> subsp. aureus derived from ATCC® 25923™*	89533	recommended for CAMP test
<i>Staphylococcus aureus</i> derived from ATCC® 6538™*	89535	
<i>Staphylococcus aureus</i> derived from ATCC® 6538P™*	89534	
<i>Staphylococcus epidermidis</i> derived from ATCC® 12228™*	89537	
<i>Streptococcus pyogenes</i> derived from ATCC® 19615™*	89538	group A



\* The ATCC Licensed Derivative Emblem, the ATCC Licensed Derivative word mark and the ATCC catalog marks are trademarks of ATCC. Liofilchem® s.r.l. is licensed to use these trademarks and to sell products derived from ATCC® cultures.

Look for the ATCC Licensed Derivative® Emblem for products derived from ATCC® cultures.

The ATCC Licensed Derivative® program was created to ensure quality and address global responsibilities for handling biological materials. A royalty fee will be charged on behalf of ATCC® for products derived from ATCC® biological material. In order to receive these products, an authorized representative from your organization must agree to the terms of the End User Agreement. It is mandatory that your organization registers for this program before we are able to process your order for lyophilized microorganism preparations containing ATCC® biological material. If you have additional questions about the program contact Liofilchem Customer Service.